

Light curve solutions of 12 eccentric *Kepler* binaries and analysis of their out-of-eclipse variability

Diana Kjurkchieva¹, Doroteya Vasileva¹ and Dinko Dimitrov²

ABSTRACT

The eccentricity, periastron angle, orbital inclination, mass ratio, stellar temperatures and relative stellar radii of 12 eclipsing eccentric binaries were determined on the basis of *Kepler* data. The analysis of their out-of-eclipse variability led to the following results: (i) KIC 10490980 exhibits rotational (spot-type) variability; (ii) Four new heartbeat stars were found: KIC 9344623 and KIC 10296163, which have wide tidally induced light humps and KIC 9119405 and KIC 9673173, which have narrow "W-shape" features; (iii) KIC 4932691 shows oscillations with approximately the 18th harmonic of the orbital period. We established that the eccentric *Kepler* binaries fall below the envelope $P(1 - e^2)^{3/2} \approx 5$ days on the period-eccentricity diagram and that there is a surprising lack of eccentric binaries with periods of 25–35 days.

Subject headings: stars: binaries: eclipsing — stars: fundamental parameters — stars: individual: (KIC 4932691, KIC 6841577, KIC 8610483, KIC 8378922, KIC 9025914, KIC 9344623, KIC 10296163, KIC 5986209, KIC 12217907, KIC 9119405, KIC 10490960, KIC 9673173)

1. Introduction

Until half a century ago the eccentric binary systems were objects mainly of the celestial mechanics. Recently they have become important targets of the modern astrophysics because their investigation provides observational tests for the theory of tidal forces and for the models of stellar structure and evolution (Fuller & Lai 2011). Eccentric binary systems have been used as probes for studying the mechanisms for circularization of the orbits, the synchronization of the stellar rotation with the orbital motion and coplanarity, the impermanent mass transfer occurring close to the periastron, and apsidal motion (Kopal 1978, Sepinsky et al. 2007, Claret & Gimenez 2010, Lajoie & Sills 2011). Despite of the recent great progress in the study of the eccentric binary stars, there remain debatable problems: the dependence of period and eccentricity and its evolutionary meaning, the empirical relations between the stellar parameters of the eccentric binaries; the existence of highly eccentric binaries with short periods, etc.

An important step in the field was made by *Kepler*. This mission's high-accuracy observations have made it possible to detect tidally induced brightening

and oscillations, which were theoretically predicted by Kumar et al. (1995). The newly discovered "heartbeat" stars represent unique laboratories for investigation of these higher - order effects (Welsh et al. 2011, Burkart et al. 2012, Thompson et al. 2012). Moreover, tidally induced pulsations could explain the driving mechanism of excitation of certain oscillation modes (Willems & Aerts 2002). The study of the Doppler boosting has also become possible (Bloemen et al. 2011).

According to the analytic model of Kumar et al. (1995) the shape of the light increase around the periastron is one-peaked for $i < 30^\circ$ but becomes two-peaked with a central dip (whose depth and width increase with i) for a steeper orbital inclination. The shapes of the tidally induced features of 16 eccentric (non-eclipsing) *Kepler* binaries discovered by Thompson et al. (2012) were varied: with a dip before brightening, with a dip after brightening, and with a distinct "W" or "M" shape. We also found tidally induced brightening for several *eclipsing* eccentric *Kepler* binaries (Kjurkchieva & Vasileva 2015a, Kjurkchieva et al. 2016, Kjurkchieva & Vasileva 2016), but with a "A" (hump) shape instead of two-peaked profile. More statistics are necessary for investigation of this newly discovered phenomenon.

This paper presents the results of our study of a sample of 12 *Kepler* binaries with narrow eclipses (widths in the range 0.006–0.036 in phase units), but

¹Department of Physics, Shumen University, 9700 Shumen, Bulgaria

²Institute of Astronomy and NAO, Bulgarian Academy of Sciences, Tsarigradsko shosse 72, 1784 Sofia, Bulgaria

Table 1: Parameters from the EB catalog: orbital period P ; *Kepler* magnitude m_K ; mean temperature T_m ; widths of the primary and secondary eclipses $w_{1,2}$ (in phase units); depths of the eclipses $d_{1,2}$ (in flux units)

Star	P [d]	m_K	T_m	w_1	w_2	d_1	d_2
KIC 4932691	18.1120792	13.627	7109	0.0134	0.0107	0.1114	0.0376
KIC 6841577	15.5375346	14.875	5478	0.0134	0.0174	0.2098	0.0171
KIC 8610483	48.7993681	15.05	5920	0.0072	0.0118	0.2783	0.1224
KIC 8378922	43.2633058	14.977	5438	0.0078	0.0157	0.3381	0.2187
KIC 9025914	11.3203214	16.756	5900	0.0146	0.02	0.2344	0.0338
KIC 9344623	14.7594793	13.822	6312	0.0223	0.019	0.3367	0.3107
KIC 10296163	9.2967644	13.211	6229	0.0262	0.0158	0.2057	0.0256
KIC 5986209	23.7379705	15.147	5345	0.0055	0.0126	0.3368	0.0868
KIC 12217907	43.2045856	14.291	5745	0.0125	0.0066	0.2047	0.0613
KIC 9119405	18.6463233	13.729	5569	0.0163	0.0208	0.2195	0.1881
KIC 10490960	5.6824111	14.3	5787	0.0327	0.0365	0.2038	0.1973
KIC 9673173	21.29474	13.731	5838	0.0136	0.0128	0.111	0.0939

still allowing acceptable light curve solutions. The aims of our investigation were (i) to obtain their orbits and physical parameters and thus to provide new data for establishment/improvement of the empirical relations between the parameters of eccentric binaries and (ii) to analyze their out-of-eclipse variability and to search for tidally induced effects.

2. Light curve solutions

Table 1 presents information for our targets from the EB catalog (Prsa et al. 2011, Slawson et al. 2011): orbital period P ; *Kepler* magnitude m_K ; mean temperature T_m ; widths of the primary and secondary eclipses $w_{1,2}$ (in phase units); and depths of the eclipses $d_{1,2}$ (in flux units). These parameters and have been obtained by an automated (pipeline) reduction of the *Kepler* data and/or preliminary ground-based photometric observations.

Long cadence (LC) data from almost all quarters are available in the *Kepler* archive for the whole sample. They reveal that besides eclipses the targets exhibit additional out-of-eclipse variability (see Section 3). To ignore this effect in the procedure of the light curve solutions, we modeled all available photometric data after appropriate phase binning.

We used the package PHOEBE (Prsa & Zwitter 2005) for modeling. The procedure is described in detail in Kjurkchieva and Vasileva (2015a). Briefly, it consists of several steps: preliminary calculation of the eccentricity e and periastron angle ω ; varying of e and ω to fit the eclipse phases; varying of the inclination i , mass ratio q , secondary temperature T_2 and potentials $\Omega_{1,2}$ (relative stellar radii $r_{1,2}$) to reproduce the whole curve; and adjusting the component temperatures around the mean target temperature T_m .

Table 2 shows the values of the orbital and stellar parameters corresponding to our light curve solutions (the periastron phase φ_{per} and the ratio of relative luminosities l_2/l_1 were not free but calculated param-

eters). The numbers in brackets in Table 2 present the parameter uncertainties obtained by PHOEBE (rounded to the greater number). The small errors of the parameters derived by our light curve solutions are natural consequence of the unique precision of the *Kepler* data.

The synthetic curves corresponding to our best light curve solutions are shown in Fig. 1 as continuous lines. The residual curves show greater scatters during the eclipse phases. Similar behavior could also be seen also for other *Kepler* binaries (Hambleton et al. 2013, Lehmann et al. 2013, Maceroni et al. 2014), especially for those with small sum of relative radii. It has been attributed to the effects of finite integration time (Kipping 2010), but we found additional reasons for these discrepancies (see Section 3.1).

The ranges of the parameter values of our targets are as follows (Table 2):

- (a) The temperatures of the stellar components correspond to spectral type from early F to late K;
- (b) The temperature differences of the components reach to around 2000 K;
- (c) The mass ratios are in the range 0.32–0.98;
- (d) The orbital inclinations are between 86.3° and 89.5° . KIC 12217907 and KIC 10296163 undergo total eclipses, while the other targets show partial eclipses;
- (e) The eccentricities of our targets are between 0.1 and 0.56.

3. Out-of-eclipse variability

All targets exhibit cyclic out-of-eclipse light variability with the orbital period (excluding KIC 4932691) of different types.

3.1. Rotational variability

The out-of-eclipse light curve of KIC 10490980 exhibits long-term modulation with amplitudes up to 0.01 mag and time scale of a hundred days (Fig. 2).




Fig. 1.— Top of each panel: the light curve of the target around the eclipses and its PHOEBE fit; bottom: the corresponding residuals.

Table 2: Parameters of the best light curve solutions: eccentricity e ; periastron angle ω ; periastron phase φ_{per} (counted from the primary minimum); inclination i ; mass ratio q ; stellar temperatures $T_{1,2}$; relative stellar radii $r_{1,2}$; ratio of relative stellar luminosities l_2/l_1

Star	e	ω [deg]	φ_{per}	i	q	T_1 [K]	T_2 [K]	r_1	r_2	l_2/l_1
KIC 4932691	0.37886(1)	181.7(7)	0.14	88.06(2)	0.393(3)	7164(48)	5473(28)	0.0436(3)	0.0153(2)	0.050
KIC 6841577	0.12280(3)	162.3(4)	0.16	88.81(3)	0.316(2)	5560(19)	3601(12)	0.0391(4)	0.0176(3)	0.046
KIC 8610483	0.3936(1)	32.9(4)	0.93	88.81(4)	0.468(4)	6158(14)	5653 (3)	0.0174(1)	0.0205(5)	0.963
KIC 8378922	0.2935(1)	112.8(2)	0.03	89.09(1)	0.573(5)	5548(20)	5356(25)	0.0229(6)	0.0234(5)	0.902
KIC 9025914	0.38029(8)	5.5(4)	0.87	88.07(6)	0.607(5)	6165(19)	4327(15)	0.0446(2)	0.0219(1)	0.052
KIC 9344623	0.18902(5)	191.5(2)	0.31	88.99(2)	0.658(5)	6395(12)	6241(12)	0.0470(2)	0.0374(1)	0.579
KIC 10296163	0.3674(1)	228.2(2)	0.29	88.53(2)	0.401(3)	6292(14)	4121(18)	0.0570(2)	0.0258(1)	0.030
KIC 5986209	0.3482(2)	55.5(2)	0.96	89.10(3)	0.339(8)	5503(25)	4868(24)	0.0188(3)	0.0144(2)	0.333
KIC 12217907	0.38819(4)	228.7(1)	0.28	89.54(3)	0.482(6)	5850(15)	4736(13)	0.0253(3)	0.0111(4)	0.078
KIC 9119405	0.44113(1)	168.3(5)	0.09	88.08(4)	0.980(1)	5549(22)	5573(24)	0.0525(4)	0.0275(3)	0.280
KIC 10490960	0.10173(7)	41.0(7)	0.88	86.32(9)	0.949(6)	5763(34)	5838(31)	0.0898(7)	0.0586(6)	0.449
KIC 9673173	0.56510(1)	0.8(4)	0.92	87.22(4)	0.752(1)	5856(12)	5683(14)	0.0436(2)	0.0158(3)	0.102

The tracing of light variability through several consecutive cycles reveals rotational (spot-type) variability (Fig. 2). Then the long-term modulation may be explained by the spot activity cycle of KIC 10490980.

3.2. New heartbeat stars

We found tidally induced light brightening with amplitude 0.0004–0.0010 mag around the periastron phase of KIC 10296163, KIC 9344623, KIC 9673173 and KIC 9119405 (Figure 3).

The features observed around the periastron of KIC 9344623 and KIC 10296163 as well as the corresponding synthetic PHOEBE features are wide and have simple "A" (hump) shapes (Fig. 3, top row).

The most interesting features at the periastron phases belong to KIC 9119405 and KIC 9673173, the targets with the greatest eccentricities from our sample. These features are relatively narrow and have complex "W-shape" profiles (Fig. 3, bottom row). The synthetic PHOEBE features reproduce the observed ones well (although not perfectly).

We found that the non-eclipsing star KIC 10873904 from the sample of Thompson et al. (2012) exhibits similar "W-shape" tidally induced brightening with amplitude of 0.001 mag (modeled by orbital parameters $e = 0.436$, $\omega = 347^\circ$ and $i = 42^\circ$).

3.3. Oscillations of KIC 4932691

The observed out-of-eclipse light curve of KIC 4932691 (Fig. 4), the target with the highest temperature, exhibits oscillations with amplitudes of 0.002–0.004 mag and a period of 1.01626 day (almost 18 times shorter than the orbital cycle). These light variations have a simple one-waved shape (Fig. 5) and could be considered as oscillations at the 18th harmonic of the orbital period, i.e. they could be attributed to the second tidally induced phenomenon predicted by Kumar et al. (1995). Most of the heartbeat targets

of Thompson et al. (2012) have oscillations that are harmonics of the orbital period, but almost all of them have higher temperatures than our targets. This implies that tidally induced oscillations are inherent to early MS stars.

3.4. Artificial light increasing at the eclipses

KIC 8610483, KIC 8378922, KIC 5986209, KIC 12217907, KIC 6841577, and KIC 9025914 reveal strange light increasing around the eclipses (Figure 6). It is more apparent at the secondary eclipse (excluding KIC 12217907) and reaches 0.0008 mag. We established that this "volcano effect" was inherent only for the targets with the narrowest eclipses ($w_{1,2} \leq 0.018$). It turned out, however, that such a feature is not present in their raw data (Fig. 7). Hence, we suppose that it is an artificial effect that is a result of the imperfect guiding of the telescope and the corresponding de-trending procedure of the *Kepler* data. This supposition explains why the synthetic PHOEBE light curves do not reproduce at all the observed light increasing around the eclipses of these targets (Fig. 6).

4. Location on the diagram period-eccentricity

One of the most interesting and debatable relations of the eccentric binaries is period-eccentricity. Different attempts have been made in the past to explain the observed trend that the binaries with longer periods have greater eccentricities, including studies of tidal action (Russell 1910); secular decrease in the stellar mass (Jeans 1924) and the effect of encounters (Walters 1932a,b). Zahn (1977) and Lecar et al. (1976) concluded that the tidal effect is especially effective for the circularization of orbits of late stars with convective envelopes.

Horrocks (1936) has derived the theoretical dependence of period and eccentricity $PM^2e^{-6}(1-e^2)^{3/2} = \text{const}$, which has been confirmed by parameters of bi-




Fig. 3.— Folded out-of-eclipse light curves of the targets with heartbeat effect.




Fig. 6.— Folded out-of-eclipse light curves of the targets with "volcano effect".




Fig. 2.— Out-of-eclipse variability of KIC 10490980 during all quarters (top) and through 10 consecutive cycles (bottom).




Fig. 7.— Out-of-eclipse light variability of KIC 8378922. Top: the automatically de-trended data with "volcano effect"; bottom: phased raw data from quarters Q5 and Q13 (shifted vertically by 0.006).




Fig. 4.— The oscillations of KIC 4932691.




Fig. 5.— The out-of-eclipse brightness of KIC 4932691 folded by period of 1.01626 d.




Fig. 8.— Period-eccentricity diagram for our 32 *Kepler* targets.

naries with periods $P \geq 5$ days. Mayor & Mermilliod (1984) studied red dwarfs in open clusters and found that the orbits of those with periods shorter than 5.7 days are circular, while binaries with longer periods have eccentric orbits. Mathieu et al. (1990) obtained similar results for 8 evolved MS binaries belonging to old open cluster M67, but determined a cut off period of around 11 days.

Duquennoy and Mayor (1991) revealed that binaries with $P \leq 10$ days have eccentricities close to zero, while those with longer periods exhibit complex (e, P) diagram. According to Raghavan et al. (2010) the period-eccentricity relation of solar-type stars shows the expected circularization for periods below 12 days (caused by tidal forces over the age of the Galaxy), followed by a roughly flat distribution for $10 \leq P \leq 1000$ days. The sample of Raghavan et al. (2010) is limited by an upper envelope $P(1 - e^2)^{3/2} = 10$ days while, the fifteen highly eccentric *Kepler* binaries studied by Dong et al. (2013) appear between the tracks of constant angular momentum corresponding to 10 days and 30 days.

The places on the period-eccentricity diagram of all 32 *Kepler* binaries we have so far studied (Kjurkchieva & Vasileva 2015a, (Kjurkchieva & Vasileva 2015b, Kjurkchieva et al. 2016, Kjurkchieva & Vasileva 2016) reveal the following trends (Figure 8): (i) There is a fast increase of the eccentricity (up to 0.57) with the period for targets with periods of 4 days $< P < 25$ days; (ii) there is a second increase of eccentricity beginning from $e = 0.1$ for periods $P \geq 35$ days; and (iii) there are no targets with periods within 25–35 days, which may mean the existence of a short stage of evolution of binaries connected with a fast change of their orbits.

We built a common (P, e) diagram for all *Kepler* eccentric binaries (Fig. 9). The targets of Dong et al. (2013) and Beck et al. (2014) fall below the envelope $P(1 - e^2)^{3/2} = 10$ days, while 14 targets (with periods below 10 days and eccentricity $e > 0.1$) from our common sample are outside of the region limited by the foregoing envelope. In fact, all our binaries fall below the envelope $P(1 - e^2)^{3/2} \approx 5$ days (Fig. 9). Most of the members of the catalog of Bulut and Demircan (2007) containing above a hundred eccentric binaries appeared on the left of the last envelope (Fig. 9).

Analysis of the common (e, P) distribution of the known eccentric binaries led us to several results.

(a) There are many short-period binaries (with periods of several days) with considerable eccentricities. Hence, a possible conclusion is that there is no sharp cut-off of the (e, P) distribution.

(b) The distribution of the eccentric binaries with

periods below 260 days reveals a roughly linear trend of the increase of eccentricity with the period (Fig. 9), which is expected from the theory of the binary evolution.

(c) There are no targets with periods of 25–35 days among all (around 180 in number) eccentric binaries shown in Fig. 9.

5. Conclusions

The main results of our study are as follows:

(1) We determined the orbital and stellar parameters of 12 eclipsing binaries with eccentric orbits from the *Kepler* archive. Their eccentricities are between 0.1 and 0.56, and most of them show partial eclipses (KIC 12217907 and KIC 10296163 have total eclipses). The temperatures of the stellar components correspond to a spectral type from early F to late K.

(2) We found out-of-eclipse light variability of different types: (i) KIC 10490980 reveals rotational (spot-type) variability. (ii) Four targets show tidally induced light features around the periastron phase - i.e. they are heartbeat stars. KIC 9344623 and KIC 10296163 have wide humps, while KIC 9119405 and KIC 9673173 (the targets with the greatest eccentricities) have narrow "W-shape" profiles. (iii) KIC 4932691 reveals oscillations with approximately the 18th harmonic of the orbital period.

(3) The eccentric *Kepler* binaries fall below the envelope $P(1 - e^2)^{3/2} \approx 5$ days on the period-eccentricity diagram. We established a lack of eccentric binaries with periods of 25–35 days.

Our investigation may be considered as expansion of the study of the heartbeat stars of Thompson et al. (2012) to the eclipsing binaries. It requires further, more in-depth analysis and interpretation in the framework of the evolutionary scenarios of binary systems.

Acknowledgements

The research was supported partly by funds of project RD 08-81/03.02.2016 of Shumen University. It used the SIMBAD database and NASA Astrophysics Data System Abstract Service. We worked with the live version of the *Kepler* EB catalog (<http://keplerebs.villanova.edu/>). The authors are grateful to the anonymous referee for the valuable recommendations and notes.

REFERENCES

- Beck, P. et al., 2014, *A & A*, 564A, 36
Bloemen, S. et al., 2011, *MNRAS*, 410, 1787

- Bulut, I., Demircan, O., 2007, MNRAS, 378, 179
- Burkart, J., Quataert, E., Arras, P., Weinberg, N.N., 2012, MNRAS, 421, 983
- Claret, A., Gimenez, A., 2010, A & A, 519A, 57
- Dong, S., Katz, B., Socrates, A., 2013, ApJ, 763L, 2
- Duquennoy, A., Mayor, M., 1991, A & A, 248, 485
- Fuller, J., Lai, D., 2012, MNRAS, 420, 3126
- Hambleton, K. et al., 2013, MNRAS, 434, 925
- Horrocks, H., 1936, MNRAS, 96, 534
- Jeans, J.H., 1924, MNRAS 85, 2
- Kipping, D., 2010, MNRAS, 408, 1758
- Kjurkchieva, D., Vasileva, D., 2015a, PASA, 32, 23
- Kjurkchieva, D., Vasileva, D., 2015b, BlgAJ, 23, 75
- Kjurkchieva, D., Vasileva, D., Dimitrov, D., 2016, Ap-SpSci, 361, 132
- Kjurkchieva, D., Vasileva, D., 2016, NewA, 48, 30
- Kopal, Z., 1978, ASSL (Astrophysics and Space Science Library) 68, Dordrecht, D. Reidel Publishing Co.
- Kumar, P., Ao, C., Quataert, E., 1995, ApJ, 449, 294
- Lajoie, C., Sills, A., 2011, ApJ, 726, 67
- Lecar, M., Wheeler, J.C., McKee, C.F., 1976, ApJ, 205, 556
- Lehmann, H., Southworth, J., Tkachenko, A., Pavlovski, K., 2013, A & A, 557A, 79
- Maceroni, C. et al., 2014, A & A, 563A, 59
- Mathieu, R., Latham, D., Griffin, R., 1990, AJ, 100, 1859
- Mayor, M., Mermilliod, J., 1984, IAUS, 105, 411
- Prsa, A., Zwitter T., 2005, ApJ, 628, 426
- Prsa, A. et al., 2011, ApJ, 141, 83
- Russell, H.N., 1910, ApJ, 31, 185
- Sepinsky, J.F., Willems B., Kalogera V., 2007, ApJ, 660, 1624
- Slawson, R. et al., 2011, AJ, 142, 160
- Thompson, S.E. et al., 2012, ApJ, 753, 86
- Walters, M.H., 1932a, MNRAS, 92, 786
- Walters, M.H., 1932b, MNRAS, 93, 28
- Welsh, W. et al., 2011, ApJS, 197, 4
- Willems, B., Aerts, C., 2002, A & A, 384, 441
- Zahn, J.-P., 1977, A & A, 57, 383




Fig. 9.— Period-eccentricity diagram of short-period eccentric binaries: pluses for the sample of Bulut and Demircan (2007); red circles for our 32 *Kepler* binaries; green triangles for the *Kepler* targets of Beck et al. (2014); blue squares for the *Kepler* binaries of Dong et al. (2013); and continuous green and violet lines describing tracks of constant angular momentum $P(1 - e^2)^{3/2}$ corresponding to 10 days and 5 days.